

University of Dundee

Obesity and asthma – do we need expensive therapies?

Anderson, William; Lipworth, Brian

Published in:
Thorax

Publication date:
2017

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Anderson, W., & Lipworth, B. (2017). Obesity and asthma – do we need expensive therapies? *Thorax*.
<http://thorax.bmj.com/content/72/2/174.responses>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Thorax

Obesity and asthma - do we need expensive therapies?

Journal:	Thorax
Manuscript ID	Draft
Article Type:	Research Letter
Date Submitted by the Author:	n/a
Complete List of Authors:	Anderson, William; University of Dundee, Scottish Centre for Respiratory Research Lipworth, Brian; University of Dundee, Scottish Centre for Respiratory Research
Keywords:	Asthma Pharmacology

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Confidential - For Review Only

Title: Obesity and asthma – do we need expensive therapies?

Authors: WJ Anderson, BJ Lipworth

Affiliation: Scottish Centre for Respiratory Research, School of Medicine,
University of Dundee, Ninewells Hospital, Dundee, DD1 9SY

Corresponding Author: Brian J Lipworth, Scottish Centre for Respiratory
Research, School of Medicine, University of Dundee,
Ninewells Hospital, Dundee, DD1 9SY

Word Count: 466

To the editor,

We read with interest the state of the art review by Umetsu on mechanisms by which obesity impacts asthma¹. He has clearly outlined the extent to which molecular targets, both within allergic (T-helper 2 [TH2]) and non-allergic mechanistic pathways of asthma, need further evaluation and drug development for obese asthmatics. This was on the basis that current standard therapies for asthma such as inhaled corticosteroids (ICS) appear to be less effective in these individuals. Whilst these novel molecular targets are clearly an exciting step forward, we believe there are several other issues that need to be addressed as well when it comes to standard asthma treatment with ICS in the obese.

Umetsu touched on altered lung mechanics in the obese stating more restriction of lung volumes². We have found previously that overweight asthmatics (body mass index [BMI] $\geq 25\text{kg/m}^2$) appear to have attenuated fractional exhaled nitric oxide (FeNO) and symptom responses as the dose of inhaled budesonide was increased up to $800\mu\text{g/day}$ compared to normal weight (BMI $< 25\text{kg/m}^2$) counterparts receiving the same ICS dose ramp³. However, there were no differences seen between responses in airway calibre as measured by forced expiratory volume in 1s (FEV₁), or indeed airway hyperresponsiveness (AHR) to methacholine. Therefore, altered corticosteroid response at a molecular level alone does not necessarily explain the equal response in these latter two outcomes. Certainly, in significantly overweight individuals their lung volumes are reduced, but we did not see this difference in our cohort, presumably reflecting a modestly increased mean BMI of 30kg/m^2 .

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

It may be, therefore, that either the ICS is not being delivered to the peripheral small airways sufficiently in obese asthmatics, in the presence of normal proximal delivery ⁴. A previous study examined the effects of bariatric surgery on measures of peripheral small airway function and asthma between asthmatic and non-asthmatic obese individuals finding that the peripheral airways are more collapsible in obese asthmatics than in their non-asthmatic counterparts⁵. Indeed small airway dysfunction itself is associated with poorer asthma control⁶ ⁷, which could feasibly have a greater impact in obese asthmatics. Therefore, strategies to improve delivery of ICS to the peripheral airways might be a more cost-effective first step for certain obese asthmatics than going directly to more expensive biological therapies to improve their asthma control. We believe further prospective study is required into the lung distribution and clinical utility of extra-fine particle ICS in obese asthmatics, as well to establish if higher than usual doses of ICS are required to achieve the same effect as in normal weight asthmatics. It goes without saying that strategies which target significant weight loss – including bariatric surgery where necessary – are likely to be even more cost-effective in the first place, as this will not only impact on improving asthma control but also on comorbidities such as diabetes and hypertension.

References

1. Umetsu DT. Mechanisms by which obesity impacts asthma. Thorax 2016.
2. King GG, Brown NJ, Diba C, et al. The effects of body weight on airway calibre. Eur Respir J 2005;**25**(5):896-901.
3. Anderson WJ, Lipworth BJ. Does body mass index influence responsiveness to inhaled corticosteroids in persistent asthma? Ann Allergy Asthma Immunol 2012;**108**(4):237-42.
4. Lipworth B, Manoharan A, Anderson W. Unlocking the quiet zone: the small airway asthma phenotype. The Lancet Respiratory medicine 2014;**2**(6):497-506.
5. Al-Alwan A, Bates JH, Chapman DG, et al. The nonallergic asthma of obesity. A matter of distal lung compliance. Am J Respir Crit Care Med 2014;**189**(12):1494-502.
6. Manoharan A, Anderson WJ, Lipworth J, et al. Small airway dysfunction is associated with poorer asthma control. Eur Respir J 2014;**44**(5):1353-5.
7. Shi Y, Aledia AS, Tatavoosian AV, et al. Relating small airways to asthma control by using impulse oscillometry in children. J Allergy Clin Immunol 2012;**129**(3):671-8.